




COVID-19 Management Protocol SGPGIMS, Lucknow

Version 1.3

June 15, 2020



Categorize based on Severity of Illness

Mild Fever, Mild URTI No dyspnoea,	Moderate Pneumonia with no signs of severe disease RR ≥ 24/ min, SPO2 ≤ 94 % on Room Air	Severe Respiratory distress requiring assisted vent. RR ≥ 30/min, SPO2 ≤ 90% on Room Air
<ul style="list-style-type: none"> Admit in Isolation Ward Contact and Droplet precautions Strict hand hygiene Tab. Hydroxychloroquine (400mg) BD on 1st day followed by 200mg 1 BD for 4 days for patients with high risk of severe disease¹. (after ECG Assessment) with Tab Azithromycin 500 mg OD x 5 days OR Tab Ivermectin 12mg OD x 3day s with Tab Doxycycline 100 mg BD x 5 days OR Tab. Favipirivir 1800mg BD on Day 1, followed by 800mg BD x 13 days Symptomatic treatment for cough and fever (bronchodilators, mucolytic, paracetamol) Monitor closely for warning signs <ul style="list-style-type: none"> Chest pain, dyspnoea Tachypnoea, cyanosis, altered mentation 	<ul style="list-style-type: none"> Admit in ICU/HDU Oxygen Support through nasal cannulae Target SpO2: 92-96% (88-92% in COPD). Awake proning as a rescue therapy. All patients should have daily 12-lead ECG Follow CRP, D-dimer & Ferritin, Fibrinogen, Procalcitonin every 48-72 hourly; CBC, KFT/LFT daily Inj. Remdesivir 200 mg IV on Day 1 followed by 100mg OD for 4 days Consider IV methylprednisolone 0.5 - 1 mg/kg or dexamethasone 0.1- 0.2 mg/kg for 3-5 days (within 48 hours of admission or if oxygen requirement is increasing and if inflammatory markers are increased) Prophylactic dose of UFH² or LMWH² (e.g., enoxaparin 40 mg per day SC) Inj. Thiamine 100 mg IV OD Inj. Vit C 1.5gm IV 6 hourly Control of co-morbid condition. Monitor for: Increased WOB, Hemodynamic instability , Increase in oxygen requirement 	<ul style="list-style-type: none"> Cautious trial of CPAP/NIV HFNC to avoid intubation Consider IV methylprednisolone 0.5 to 1 mg/kg dexamethasone 0.1- 0.2 mg/kg for 5-7 days if not already given Therapeutic dose of UFH or LMWH (e.g., enoxaparin 40 mg per day SC) after excluding coagulopathy or thrombocytopenia or high risk of bleeding³ Inj. Thiamine 100 mg IV OD Inj. Vit C 1.5gm IV 6 hourly Mechanical ventilation if unable to maintain saturation, increased work of breathing or development of hemodynamic instability <ul style="list-style-type: none"> Conventional ARDS Net strategy Proning, recruitment manoeuvres Management of septic shock as per SSC guidelines and local antibiotic policy Convalescent Plasma (Under Trial Setting) or rescue therapy on compassionate grounds ** Tocilizumab (Off label, Individualise),

Testing
While attending suspect case as per above protocol based on clinical assessment, testing shall be resorted to and if negative—manage in Non-Covid facility according to clinical diagnosis

Discharge
After clinical improvement, discharge according to state discharge policy

1. High risk patients for Severe Disease
 - Age > 60 years
 - HTN, Diabetets Mellitus and other immunocompromising conditions.
 - Chronic lung, kidney or liver disease
 - Cerebrovascular disease
 - Obesity BMI > 25 kg/m²

2. LMWH: Low Molecular Weight Heparin; if no contraindication or high risk of bleeding; UFH: Unfractionated Heparin
 3. Risk of Bleeding: Use validated score for assessing bleeding risk (e.g. HAS-BLED Score), Use D-Dimer and SIC for further risk stratification (SIC score ≥ 24 portends high thrombotic risk)
 * Apply Emergency Severity Index (ESI): ESI: 1-2—Unstable, ESI: 3—Borderline, ESI: 4-5—Stable
 # Nasopharyngeal/Oropharyngeal Swab
 ** Informed consent mandatory before use of off label drugs.